arXiv:1201.0663v2 [quant-ph] 5 Jul 2012

Motion-generated quantum gates and entanglement resonance

David Edward Bruschi,[1] Andrzej Dragan,[1, 2] Antony R. Lee,[1] Ivette Fuentes,[1, *] and Jorma Louko[1]
[1]*School of Mathematical Sciences, University of Nottingham,*
University Park, Nottingham NG7 2RD, United Kingdom
[2]*Institute of Theoretical Physics, University of Warsaw, Hoża 69, 00-049 Warsaw, Poland*
(Dated: July 9, 2012)

The motion of a cavity in spacetime generates quantum gates via entanglement between cavity modes. While entanglement produced by uniform acceleration is small, we show analytically that two-mode quantum gates with an observable amount of entanglement can be generated by periodic repetition of any non-uniform trajectory segment. A resonance occurs when the frequency associated with the duration of the repeated segment equals half the frequency sum of any two oddly separated cavity modes, leading to an entanglement accumulation that grows linearly in the number of segment repetitions.

Introduction. — Understanding how motion and gravity affect entanglement is a key feature in the implementation of new quantum information technologies, including quantum cryptography and teleportation, in space-based scenarios that are currently under investigation [1]. Quantifying entanglement in situations where motion or gravitation have a significant role can also provide guidance for theories about the microscopic structure of spacetime, via the Hawking-Unruh effect and its connections to thermodynamics and statistical mechanics [2, 3]. Recent work in relativistic quantum information has shown that non-uniform motion of a cavity creates entanglement between the cavity field modes [4, 5]. In this Letter we show that by repeating any trajectory segment periodically it is possible to generate a resonance creating higher amounts of entanglement for small cavities and large accelerations or large cavities and small accelerations. Non-uniform motion induces transformations on the states which are equivalent to two-mode squeezing operations upto local unitaries. These transformations play the role of two-mode quantum gates in continuous variables systems[6].

Finding suitable ways to store and process information in a quantum and relativistic setting is a main goal in the field of relativistic quantum information. Moving cavities are good candidates to store information [7–9] since confined fields can be realized experimentally and observers can directly access their states by means of local operations. When a cavity is accelerated for a finite time the cavity modes are affected by the motion. A mismatch between the vacua at different times gives rise to the creation of particles which populate the modes [10]. The initial and final modes are related through Bogoliubov transformations which mix all frequency modes [4, 7] generating entanglement [4, 5]. Quantum correlations are created between all modes; however, higher degrees of entanglement are produced between oddly separated modes. As a consequence, maximally entangled states of modes confined in two different cavities degraded when the cavities undergo non-uniform motion [7].

We show that two-mode squeezed states can be produced in a single cavity by the periodic repetition of any trajectory segment. Entanglement resonances occur when the frequency associated to the segment travel time is equal to the sum of two oddly separated cavity mode frequencies. We show analytically that for *any* couple of oddly separated modes it is possible to find a segment travel time where the entanglement between such modes increases linearly with the number of repetitions. These resonances appear independently of the details of the trajectory though the amount of entanglement generated does depend on the trajectory itself.

As a concrete example we present a travel scenario which allows for simple analytical expressions. The travel segment which we repeat starts with a period of acceleration followed by inertial coasting. The cavity then accelerates in the same or opposite direction for a second period of time and finally undergoes a second period of inertial coasting. We find the conditions for the resonances to occur and investigate how the magnitude and the direction of the accelerations affect the final entanglement generated. In the special case where there is no inertial coasting and the accelerations alternate in direction, our sample trajectory reduces to the oscillatory motion that is often considered in the dynamical Casimir effect literature [11].

Setup. — We consider a real scalar field ϕ of mass m contained within a cavity in $(1+1)$-dimensional spacetime, with Dirichlet boundary conditions at the walls. The massless field can be treated as a special case of our study and the effect of additional transverse dimensions can be included as a positive contribution to m. The cavity follows a worldtube which is composed of periods of inertial and uniformly accelerated motion. We will start by describing the field within the cavity during such periods as seen by a co-moving observer.

During periods of inertial motion, we use Minkwoski coordinates (t, x) to describe our system. The cavity walls are placed at $x = x_A$ and $x = x_B$ where $0 < x_A < x_B$, so that $L = x_B - x_A$ is the length of the cavity. The positive frequency mode functions with respect to the time translation Killing vector ∂_t are:

$$\phi_k^M(t,x) = \frac{1}{\sqrt{\omega_k L}} \sin\left[\frac{k\pi}{L}(x - x_A)\right] e^{-i\omega_k t} \ ,$$
$$\omega_k = \sqrt{(k\pi/L)^2 + m^2} \ , \tag{1}$$

where $k \in \mathbb{N}$ (we set $c = \hbar = 1$).

We employ Rindler coordinates (τ, χ) to describe the field during periods of uniformly accelerated motion. Rindler coordinates are related to Minkowski coordinates through the transformations $\chi = \sqrt{x^2 - t^2}$ and $\tau = (1/a)\,\mathrm{atanh}(t/x)$, where a is a positive constant. An uniformly accelerated observer comoving with the cavity follows the boost Killing vector field $\partial_\eta = x\partial_t + t\partial_x$ which in Rindler coordinates takes the form $\partial_\eta = (1/a)\partial_\tau$. The cavity walls are placed at $\chi = x_A$ and $\chi = x_B$ and the proper time and acceleration at the centre of the cavity are given by τ and $a = 2/(x_B + x_A)$, respectively. The solutions to the Klein-Gordon equation that are of positive frequency with respect to ∂_η are

$$\begin{aligned}\phi_k^R(\tau,\chi) =& N_k \left[I_{i\Omega_k/a}(m\chi) I_{-i\Omega_k/a}(m x_A) \right. \\ & \left. - I_{-i\Omega_k/a}(m\chi) I_{i\Omega_k/a}(m x_A) \right] e^{-i\Omega_k \tau} \ ,\end{aligned} \tag{2}$$

where I_ν are modified Bessel functions of the first kind [12] and N_k are normalisation constants. In the massless case the mode functions and the frequencies reduce to simple expressions (see [7]).

The quantized field operators are given by $\phi^M = \sum_k \left(\phi_k^M a_k + \text{h.c.}\right)$, and $\phi^R = \sum_k \left(\phi_k^R A_k + \text{h.c.}\right)$ during periods of inertial and accelerated motion, respectively. The Minkowski and Rindler annihilation and creation operators obey the standard commutation relations $\left[a_k, a_l^\dagger\right] = \delta_{kl}$ and $\left[A_k, A_l^\dagger\right] = \delta_{kl}$. The Minkowski frequencies, $\omega_k > 0$, are determined by the Dirichlet boundary condition $\phi_k^M(t, x_B) = 0$ and given by (1), while Rindler frequencies, $\Omega_k > 0$, are determined by $\phi_k^R(\tau, x_B) = 0$.

We will work in the covariant matrix formalism which is applicable to systems consisting of a discrete number of bosonic modes as long as the analysis is restricted to gaussian states. In this framework the state of the system is entirely described by its first and second moments [13, 14]. The evolution of the state is given by a similarity transformation $S^{Tp}\Delta S$ where S is the symplectic representation of the evolution and Δ is the covariance matrix encoding all information pertinent to the state. Tp denotes matrix transposition.

Travel scenario techniques. — Changes from inertial to accelerated motion and vice versa are implemented by the action of Bogolubov transformations. Consider that at $t = 0$ a cavity initially at rest begins to accelerate. The inertial and accelerated cavity modes are related by the Bogolubov transformation $\phi_k^R = \sum_n \left(\alpha_{kn}\phi_n^M + \beta_{kn}\phi_n^{M*}\right)$, where the star denotes complex conjugation and the Bogolubov coefficients α_{kl} and β_{kl} can be evaluated by taking Klein-Gordon inner products of the two sets of modes at $t = 0$ [10, 15, 16]. In the low acceleration regime we can expand the coefficients in a Maclaurin series as $\alpha = \mathbb{I} + h\alpha^{(1)} + O(h^2)$ and $\beta = h\beta^{(1)} + O(h^2)$ where $h = aL$ is a small dimensionless expansion parameter [7, 8]. The coefficients can be computed using uniform asymptotic expansions of Bessel functions [12, 17].

During the periods of inertial or accelerated motion the modes undergo free evolution which induces phase rotations on the state of the form $U(\tau) = \bigoplus_{i=1}^N R(\theta_i)$ where N is the number of bosonic modes considered and R is the rotation matrix,

$$R(\theta_k) := \begin{pmatrix} \cos\theta_k & \sin\theta_k \\ -\sin\theta_k & \cos\theta_k \end{pmatrix} . \tag{3}$$

The angles are given by $\theta_k = \omega_k t$ during coasting periods and $\theta_k = \Omega_k \tau$ during acceleration. The Rindler and Minkowski frequencies coincide to first order ($\Omega_k(h) \to \omega_k a$ as $h \to 0$).

By composing these basic transformations we construct our trajectores. The field modes of a cavity initially at rest and the modes after any travel scenario are related through general Bogolubov transformations. In the covariant matrix formalism the transformations are represented the symplectic matrix

$$\mathcal{S} = \begin{pmatrix} s_{11} & s_{12} & \dots \\ s_{21} & s_{11} & \dots \\ \vdots & \vdots & \ddots \end{pmatrix} , \tag{4}$$

where the 2×2 matrices $s_{kk'}$ take the form

$$s_{kk'} = \begin{pmatrix} \Re(A_{kk'} - B_{kk'}) & \Im(A_{kk'} + B_{kk'}) \\ -\Im(A_{kk'} - B_{kk'}) & \Re(A_{kk'} + B_{kk'}) \end{pmatrix} , \tag{5}$$

where $A_{kk'}$ and $B_{kk'}$ are Bogolubov coefficients associated with the whole trajectory [5]. In the case of a cavity initially at rest that begins to uniformly accelerate at $t = 0$ the symplectic matrix, which we denote by $\mathcal{V}$, corresponds to $A_{kk'} = \alpha_{kk'}$ and $B_{kk'} = \beta_{kk'}$. When the Bogolubov coefficients $\beta_{kk'}$ are non-vanishing there is particle creation and the cavity modes become entangled according to a co-moving observer. The basic building block trajectory which corresponds to inertial-uniformly accelerated-inertial motion, is implemented by the action of the symplectic matrix $S_B = \mathcal{V}^{-1}(h) U(\tau) \mathcal{V}(h)$. The entanglement generated between the cavity modes after a single basic building block trajectory has been analyzed in [5] and found to be very small.

More general travel scenarios are constructed by composing basic building block trajectories. Here we are interested in constructing trajectory segments which will then be repeated to generate resonances. A segment contains any number of different basic building blocks and could be for example, a return voyage to Alpha-Centauri [7]. We consider the cavity to be initially in the vacuum state which satisfies $a_k|0\rangle = 0 \ \forall k$ and is represented by the identity matrix $\Delta = \mathbb{I}$. This choice is convenient since the state is gaussian and separable (not entangled) in the mode number degree of freedom k. In this case, it is possible to calculate the reduced state of two field modes k and k' after the cavity has followed a general trajectory segment as long as the product of the acceleration and the cavity length remains small. We assume the different accelerations involved in the segment to be of the form $a_i = s_i a$ where a is the largest acceleration and $s_i < 1$ a real number. The Bogolubov coefficients can be expanded to first order in $h = aL$ as $A_{kk'} = G_k \delta_{kk'} + A^{(1)}_{kk'}$ and $B_{kk'} = B^{(1)}_{kk'}$ where the superscript (1) denotes the first order correction and $G_k = e^{i\omega_k T}$ are the phases acquired by the state during periods of free evolution. T denotes the *total* proper time of the segment. When the paramater h is small and the initial state is the vacuum, the reduced state of two modes after a segment trajectory is given by $\sigma_1 = S^{Tp}_{kk'} S_{kk'}$ and by $\sigma_N = (S^N_{kk'})^{Tp} S^N_{kk'}$ after the segment has been repeated N times. The matrix $S_{kk'}$ takes the form

$$S_{kk'} = \begin{pmatrix} s_{kk} & s_{kk'} \\ s_{k'k} & s_{k'k'}, \end{pmatrix} \qquad (6)$$

Entanglement resonance and two-mode quantum gates — Our aim is to quantify the entanglement of the reduced state σ_N. We find that, to first order in h, the final reduced state of two modes is separable for evenly separated modes but entangled and pure for oddly separated modes since $\mathrm{Tr}(\sigma_1) = 1 + \mathcal{O}(h^2)$ [5]. The mixedness between k, k' and all other modes contributes only at second order $\mathcal{O}(h^2)$ [5]. The entanglement of a bipartite system in a pure state is quantified by the von Neumann entropy; however, this measure is not suitable in our perturbative regime because the first order contribution in the expansion of the entropy is of the form $\mathcal{E}_{VN} \sim h \ln h$ which cannot appear as a term in a power series expansion. Fortunately, lower bounds to the entanglement generated in the system can be found by calculating the smallest positive symplectic eigenvalue $\tilde{\nu}_N$ of the partial transposed state $\tilde{\sigma}_N = P\sigma_N P$ [14] where $P = \mathrm{diag}(1,1,1,-1)$. This symplectic eigenvalue is found by diagonalizing the matrix $M = i\Omega\tilde{\sigma}_N$ where Ω is the antisymmetric symplectic form which, written in terms of Pauli matrices σ_y, takes the form $\Omega = -\sigma_y \oplus \sigma_y$. One can show that $\tilde{\nu}_N$ has a well defined power series expansion in h. There is a class of entanglement measures, including the logarithmic negativity, based on the *positive partial transpose criterion* [18] which are monotonic functions of $\tilde{\nu}_N$ and therefore, also have a suitable power series expansion.

The perturbative expansion of the symplectic eigenvalue is given by $\tilde{\nu}_N = 1 - \tilde{\nu}^{(1)}_N$ where $0 \leq \tilde{\nu}^{(1)}_N \propto h$ is the first order correction. We find that when the commutator $[S^{Tp}_{kk'}, S_{kk'}]$ vanishes the partial transposed state after N segment repetitions is equal to the N-th power of the partial transposed state after a single repetition of the segment, i.e. $\tilde{\sigma}_N = \tilde{\sigma}^N_1$. After simple calculations it is possible to show that to first order, when the commutator vanishes, the partial transpose state grows linearly with the number of repetitions $\tilde{\sigma}_N = \mathbb{1} + \tilde{\sigma}^{(1)}_N = \mathbb{1} + N\tilde{\sigma}_1$. After diagonalizing $\tilde{\sigma}_1$ and $\tilde{\sigma}_N$ we find that the corrections to the eigenvalues satisfy the relation $\tilde{\nu}^{(1)}_N = N\tilde{\nu}^{(1)}_1$. Any entanglement measure $E(\tilde{\nu}_N)$ that is a function of the symplectic eigenvalue $\tilde{\nu}_N$ will have a power expansion with $\tilde{\nu}^{(1)}_N$ as a leading term. In particular, when $\tilde{\nu}^{(1)}_N = \tilde{\nu}^{(1)}_1$, entanglement grows linearly

$$E(\tilde{\nu}_N) \sim \tilde{\nu}^{(1)}_N \sim N\tilde{\nu}^{(1)}_1 \sim NE(\tilde{\nu}_1) \qquad (7)$$

and therefore $[S^{Tp}_{kk'}, S_{kk'}] = 0$ is a *resonance condition.* In particular, the logarithmic negativity at resonance is given by $E_{\mathcal{N}} = N\tilde{\nu}_1$. Note that the condition $N\tilde{\nu}^{(1)}_1 \ll 1$ must be satisfied to remain within the perturbative regime [5]. The state we consider is pure, bipartite and symmetric. It is well known in the covariant matrix formalism that in this case the state is equivalent to a two mode squeezed state upto local operations which do not affect the entanglement [14]. Operations that produce two-mode squeezed states are used as two-mode quantum gates in continuos variable systems. Using this fact one can show that $\tilde{\nu}^{(1)}_N \sim B^{(1)}_{kk'}$ where $B^{(1)}_{kk'}$ is the first order correction to the beta coefficients $B_{kk'}$ of the segment [5]. Therefore, a lower bound to the entanglement generated at resonance after N segment repetitions will be given by the logarithmic negativity $E_{\mathcal{N}} = NB^{(1)}_{kk'}$. We find that the commutator for an arbitrary segment to first order in h, is

$$\left[S^{Tp}_{kk'}, S_{kk'}\right] = \begin{pmatrix} 0 & C \\ C^T & 0 \end{pmatrix}, C = 2\begin{pmatrix} C_{11} & C_{12} \\ C_{12} & C_{22} \end{pmatrix} \qquad (8)$$

where

$$\begin{aligned} C_{11} &= \Big[(\cos(\omega_k T) - \cos(\omega_{k'} T))\Re(B^{(1)}_{kk'}) \\ &\quad + (\sin(\omega_k T) + \sin(\omega_{k'} T))\Im(B^{(1)}_{kk'})\Big] \\ C_{12} &= \Big[(-\cos(\omega_k T) + \cos(\omega_{k'} T))\Im(B^{(1)}_{kk'}) \\ &\quad + (\sin(\omega_k T) + \sin(\omega_{k'} T))\Re(B^{(1)}_{kk'})\Big]. \end{aligned} \qquad (9)$$

Since the Minkowski and Rindler frequencies coincide to order h^1, the coefficients C_{ij} depend only on the Minkowski frequencies ω_k. We find that the commutator vanishes when the following equation is satisfied

$$B^{(1)}_{kk'} \left[G^*_k - G_{k'}\right] = 0. \qquad (10)$$

Therefore, resonances occur when $B^{(1)}_{kk'} \neq 0$ and the total proper time T takes discrete values

$$T_n = \frac{2n\pi}{\omega_k + \omega_{k'}}. \qquad (11)$$

We emphasize that the value of T_n does not depend on the details of the travel scenario; however, the total amount of entanglement generated $E_{\mathcal{N}} = NB^{(1)}_{kk'}$ does depend on the specifics of the trajectory thorough the Bogolubov coefficient. It is also possible to calculate the average number of excitations $\langle N_k \rangle$ in each cavity mode while on resonance. We find with the result is also proportional to $\tilde{\nu}^{(1)}_N$.

Sample travel scenario. — We now specialize to our sample travel scenario which corresponds to a cavity which is initially inertial, travels with proper acceleration $a = h/L$ for proper time τ, coasts for proper time t, travels with proper acceleration (in either the same or opposite direction) $a' = h'/L$ for proper time τ and finally coasts for proper time t. The total proper time of the segment is $T_n = 2(\tau + t)$ and the symplectic transformation is

$$\begin{aligned} S_{kk'} =& U_{kk'}(t)\mathcal{V}^{-1}_{kk'}(h')U_{kk'}(\tau)\mathcal{V}_{kk'}(h')\times \\ & U_{kk'}(t)\mathcal{V}^{-1}_{kk'}(h)U_{kk'}(\tau)\mathcal{V}_{kk'}(h). \end{aligned} \tag{12}$$

In this case, we find an analytical expression for the first order correction to the beta Bogolubov coefficient,

$$|B^{(1)}_{kk'}| = c_{kk'}|1 - g^*_k g^*_{k'}||1 + \epsilon y g^*_k g^*_{k'} f^*_k f^*_{k'}|h \tag{13}$$

where the phases are given by $g_k = \exp(i\omega_k\tau)$ and $f_k = \exp(i\omega_k t)$ and we chose $a' = \epsilon y a$ $(y > 0)$. The parameter ϵ is equal to 1 when the acceleration of the second period has same the direction as a and -1 when the direction of acceleration changes. $c_{kk'} = \frac{\sqrt{kk'}(1-(-1)^{k-k'})}{\pi^2(k+k')^3}$ are coefficients which vanish for evenly separated modes. Substituting Eq. (11) in Eq. (13) we find that the logarithmic negativity at resonant times is,

$$E_{\mathcal{N}} = NC_{kk'}|(1-(-1)^n e^{i(\omega_k+\omega'_k)t})(1+(-1)^n\epsilon y)|h. \tag{14}$$

Note that the logarithmic negativity vanishes when n is even and the time of coasting is $t = 2\pi m/(\omega_k + \omega'_k)$ and when n is odd and $t = (2m+1)\pi/(\omega_k + \omega'_k)$ with $m \in \mathbb{N}$. In the case the accelerations have the same magnitude $(y = 1)$ maximal amounts of entanglement are generated when n is even and $t = \pi(2m+1)/(\omega_k + \omega'_k)$ in the case when the accelerations have the same direction $(\epsilon = 1)$ and when n is odd and $t = 2\pi m/(\omega_k + \omega'_k)$ when the accelerations alternate direction $(\epsilon = -1)$. This behavior is shown in Fig. 1 where we plot the symplectic eigenvalue $\tilde{\nu}^{(1)}_N$ after $N = 5$ segment repetitions as a function of the proper time of acceleration τ and the time of coasting t. We considered a cavity length of $L = 1$, massless modes $k = 1$ and $k' = 2$ and accelerations $a = a' = 10^{-4}$.

Interestingly, the special case of alternating acceleration directions $(\epsilon = -1)$ and no coasting $(t = 0)$ corresponds to the standard dynamical Casimir setting where the cavity oscillates periodically as a whole. A resonant enhancement of particle creation occurs in the dynamical Casimir effect [11] which was recently demonstrated in the laboratory in a superconducting circuit consisting of a coplanar transmission line with a tunable electrical length which produces an effective moving boundary [19].



FIG. 1. The correction to the symplectic eigenvalue $\tilde{\nu}^{(1)}_N$ after $N = 5$ segment repetitions as a function of the proper time of acceleration τ and the time of coasting t. We considered a cavity length of $L = 1$, massless modes $k = 1$ and $k' = 2$ and accelerations $a = a' = 10^{-4}$.

Discussion. — We have shown that non-uniform motion can generate two-mode quantum gates which produce observable amounts of entanglement. Finding ways to create significant amounts of entanglement in relativistic settings is of great interest since entanglement is necessary for quantum communications and information processing [20]. Recent studies in relativistic quantum information show that small amounts of mode entanglement are created when a cavity undergoes non-uniform motion. [7, 8]. We show that particle creation and bipartite mode entanglement can be linearly enhanced by repeating any travel segment periodically. Via the equivalence principle our results suggest that fluctuations of a gravitational field can produce entanglement. For example, consider a small cavity containing a bosonic field in its vacuum state free falling in the presence of a gravitational field [21]. Entanglement between the modes is generated by suddenly holding the cavity at a fixed position against the action of the gravitational field. If the cavity's position changes periodically or the gravitational field fluctuates, the entanglement can be enhanced.

We thank Gerardo Adesso, Nicolai Friis and Davide Girolami for interesting discussions and useful comments. I. F. and A. D. thank EPSRC [CAF Grant EP/G00496X/2] for financial support. A. D. was supported in part by the Polish Ministry of Education. J. L. was supported in part by STFC (UK).

---

[*] Previously known as Fuentes-Guridi and Fuentes-Schuller.

[1] P. Villoresi et al, New Journal of Physics **10**, 033038 (2008).

[2] S. Hawking, Commun.Math.Phys. **43**, 199 (1975).

[3] W. Unruh, Phys.Rev.D **14**, 870 (1976).

[4] N. Friis, D. E. Bruschi, J. Louko, and I. Fuentes, Phys. Rev. D **85**, 081701 (2012).

[5] N. Friis and I. Fuentes, "Motion generates entanglement," (2012), arXiv:1201.0549v3 [quant-ph].

[6] C. Weedbrook, S. Pirandola, R. García-Patrón, N. J. Cerf, T. C. Ralph, J. H. Shapiro, and S. Lloyd, Rev. Mod. Phys. **84**, 621 (2012).

[7] D. E. Bruschi, I. Fuentes, and J. Louko, Phys. Rev. D **85**, 061701 (2012).

[8] N. Friis, A. R. Lee, D. E. Bruschi, and J. Louko, Phys. Rev. D **85**, 025012 (2012).

[9] T. G. Downes, I. Fuentes, and T. C. Ralph, Phys. Rev. Lett. **106**, 210502 (2011).

[10] N. Birrell and P. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).

[11] V. Dodonov, A. Klimov, and V. Manko, Physics Letters A **149**, 225 (1990).

[12] Digital Library of Mathematical Functions. Release date 2012–03–23. National Institute of Standards and Technology, http://dlmf.nist.gov/10.25 .

[13] S. L. Braunstein and P. van Loock, Rev. Mod. Phys. **77**, 513 (2005).

[14] J. Laurat et. al., Journal of Optics B: Quantum and Semiclassical Optics **7**, S577 (2005).

[15] A. Fabbri and J. Navarro-Salas, *Modeling black hole evaporation* (2005).

[16] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, Rev. Mod. Phys. **80**, 787 (2008).

[17] T. M. Dunster, SIAM J. Math. Anal. **21**, 9951018 (1990), note a misprint in formula (4.15).

[18] A. Peres, Phys. Rev. Lett. **77**, 1413 (1996).

[19] C. M. Wilson, G. Johansson, A. Pourkabirian, J. R. Johansson, T. Duty, F. Nori, and P. Delsing, Nature **479**, 18 (2010).

[20] R. Schützhold, G. Schaller, and D. Habs, Phys. Rev. Lett. **100**, 091301 (2008).

[21] D. Kothawala, "Box of ideal gas in free fall," arXiv:1108.6301 [gr-qc].